




Booster[®] Concentrate

A patented highly purified distilled monoglyceride

HEALx Booster[®] Concentrate is a patented highly purified distilled monoglyceride. It is used in HEALx Soother Plus[®] and Booster[®] products, and may be administered orally in the concentrated form.

TECHNICAL INFORMATION

It is important to know what a monoglyceride like monolaurin (Booster[®] Concentrate) is. For visualization, a monoglyceride is composed of glycerin joined with one fatty acid (see drawing below). In this illustration, “sn” represents the glycerol carbon position, “n” the length and “R” the kind of fatty acid attached to glycerol.



A monoglyceride (MG) lipid, of which there are two kinds, will have only one fatty acid attached to the sn1(3)- or the sn2-position of glycerol. The fatty acids in the sn1 and sn3 positions are similar since they occupy a position at either end of the glycerol molecule. However, not all monoglycerides are the same since the fatty acid can occupy the end position sn1(3) of glycerol or the middle sn2 position. Their structural differences need to be emphasized. It is important therefore to remember these structural sn designations because they reflect the different biological metabolism of the two monoglycerides. Where there are two fatty acids attached to glycerin, the lipid is known as a diglyceride (DG). If three fatty acids are attached, the lipid is known as a triglyceride (TG) or simply as a fat.

The antiviral, antibacterial, and antiprotozoal properties of lauric acid and monolaurin have been recognized for nearly three decades by only a small number of researchers: their work, however, has resulted in 100 or more research papers and numerous U.S. and foreign patents. While nontoxic and approved as a GRAS direct food additive by the FDA, Booster[®] Concentrate adversely affects bacteria, yeast, fungi, protozoa and enveloped viruses.

Researchers found that the properties that determine the anti-infective action of lipids are related to their structure: e.g., free fatty acids and monoglycerides. While the monoglycerides are active, diglycerides and triglycerides (fats) are inactive. Of the saturated fatty acids, lauric acid has greater antiviral activity than caprylic acid (C-8), capric acid (C-10), or myristic acid (C-14).

Fatty acids and monoglycerides produce their killing/inactivating effects by several mechanisms. An early postulated mechanism was the perturbing of the plasma membrane lipid bilayer.



The antiviral action attributed to monolaurin is that of fluidizing the structure in the envelope of the virus, causing the disintegration of the microbial membrane. More recent studies indicate that one antimicrobial effect in bacteria is related to monolaurin's interference with signal transduction/toxin formation (Projan et al 1994). Another antimicrobial effect in viruses is due to lauric acid's interference with virus assembly and viral maturation (Homung et al 1994). The third mode of action may be on the immune system itself (Witcher et al 1996).

MICROORGANISMS INACTIVATED BY BOOSTER[®] CONCENTRATE IN LABORATORY STUDIES

VIRUSES

- HIV or HIV-1, -6
- Herpes simplex virus-i (HSV-1 & 2)
- Measles virus
- Epstein-Barr virus (EBV)
- Influenza virus
- Leukemia virus
- Semliki forest virus
- Human papillomavirus (HPV)
- Visna virus
- Vesicular stomatitis virus (VSV)
- Rubella virus
- Respiratory syncytial virus
- Dengue virus (Type 1-4)
- Cytomegalovirus (CMV)
- Lymphocytic choriomeningitis
- Pneumovirus

BACTERIA

Gram-positive organisms

- *Bacillus anthracis* (anthrax)
- *Listeria monocytogenes*
- *Staphylococcus aureus*
- Groups A, B, F & G streptococci
- *Streptococcus agalactiae*
- *Mycobacterium* sp.
- *Clostridium perfringens*

Gram-negative organisms

- *Chlamydia pneumoniae*
- *Neisseria gonorrhoeae*
- *Helicobacter pylorus*
- *Mycoplasma pneumoniae*
- *Vibrio parahaemolyticus*

OTHERS IF USED CONCURRENTLY WITH A CHELATOR

Yeasts, fungi and molds

- *Aspergillus niger*
- *Saccharomyces cerevisiae*
- Ringworm or tinea (*Trichophyton*)
- *Malassezia* sp.
- *Penicillium citrinum*
- *Candida utilis*
- A number of protozoa-like *Giardia lamblia*

ANTIVIRAL EFFECTS

Hierholzer and Kabara (1982) first reported the antiviral activity of the monoglyceride of lauric acid (monolaurin) on viruses that affect humans. They showed virucidal effects of monolaurin on enveloped RNA and DNA viruses. This work was done at the Centers for Disease Control of the U.S. Public Health Service. This study was carried out using selected virus prototypes or recognized representative strains of enveloped human viruses. All these viruses have a lipid membrane. The presence of a lipid membrane on viruses makes them especially vulnerable to lauric acid and its derivative monolaurin. These initial findings from the Centers for Disease Control (CDC) have been confirmed by many other investigators.

Research has shown that enveloped viruses are inactivated by added fatty acids and monoglycerides in both human and bovine milk (Isaacs et al 1991). Others (Isaacs et al 1986, 1991; Thormar, et al 1987) have confirmed original statements concerning the effectiveness of monolaurin.

Some of the viruses inactivated by these lipids are the measles virus, herpes simplex virus (HSV-1 and -2), herpes family members (HIV, hepatitis C, vesicular, stomatitis virus [VSV], visna virus, and cytomegalovirus [CMV]). Many of the pathogenic organisms reported to be inactivated by these antimicrobial lipids are those known to be responsible for opportunistic infections in HIV-positive individuals. For example, concurrent infection with cytomegalovirus is recognized as a serious complication for HIV-positive individuals (Macallan et al 1993).

Thus, it would appear imperative to investigate the practical aspects and the potential benefit of a nutritional supplement such as Booster® Concentrate for microbial-infected individuals. Until now few nutritionists in the mainstream nutrition community seem to have recognized the added benefit of antimicrobial lipids in the support of infected patients. These antimicrobial fatty acids and their derivatives are essentially nontoxic to man. According to the published research, lauric acid is one of the best "inactivating" fatty acids, and its monoglyceride is even more effective than the fatty acid alone (Kabara 1978, Sands et al 1979, Fletcher et al 1985, Kabara 1985).

It should be emphasized that lauric acid cannot be taken orally because it is severally irritating. Booster® Concentrate, on the other hand is a derivative of lauric acid chemically bonded to glycerin to form monolaurin and can be taken orally without any problem.

ANTIBACTERIAL EFFECTS

The potentially pathogenic bacteria inactivated by monolaurin include *Listeria monocytogenes*, *Staphylococcus aureus*, *Streptococcus agalactiae*, Groups A, streptococci-gram-positive organisms, and some gram-negative organisms (*Vibrio parahaemolyticus* and *Helicobacter pylori*).

Decreased growth of *Staphylococcus aureus* and decreased production of toxic shock syndrome toxin-I were shown with monolaurin (Holland et al 1994). Monolaurin was 5000 times more inhibitory against *Listeria monocytogenes* than ethanol (Oh & Marshall 1993). In vitro monolaurin rapidly inactivated *Helicobacter pylori*. Of greater significance there appears to be very little development of resistance of

the organisms to the bactericidal effects (Petschow et al 1996) of these natural antimicrobials.

A number of fungi, yeast and protozoa are also inactivated or killed by monolaurin. The fungi include several species of ringworm (Isaacs et al 1991). The yeast reported to be affected is *Candida albicans* (Isaacs et al 1991). The protozoan parasite *Giardia lamblia* is killed by monoglycerides from hydrolyzed human milk (Hemell et al 1986, Reiner et al 1986, Crouch et al 1991, Isaacs et al 1991).

Chlamydia trachomatis is inactivated by monolaurin (Bergsson et al 1998). Hydrogels containing monocaprin/monolaurin are potent in vitro inactivators of sexually transmitted viruses, such as HSV-2 and HIV-1, as well as bacteria, such as *Neisserian gonorrhoea* (Thormar et al 1999).

Monolaurin does not appear to have an adverse effect on desirable gut bacteria but rather on only potentially pathogenic microorganisms. For example, Isaacs et al (1991) reported no inactivation of the common *Esherichia coli* or *Salmonella enteritidis* by monolaurin, but major inactivation of *Hemophilus influenzae*, *Staphylococcus epidermis* and Group B gram-positive *Streptococcus*.

THE PROBLEM OF ANTIBIOTICS

The phenomenal rate of prescriptions dispensed for antibiotic use, and to a lesser extent antiviral use, has grown exponentially in the past several decades. An antibiotic has limited specificity and generally does not recognize "good" bacteria (often referred to as probiotics or for life) from "bad" bacteria (meaning those bacteria that may cause disease). Antibiotics try to destroy all bacteria and are usually unsuccessful.

More antibiotic therapy may start perpetuating a chronic illness. The cycle of antibiotic therapy may go on for months and months, and repetitive indiscriminate use of antibiotics destroys weak bacteria and sets up the stage for the more virulent bacteria to survive (as in survival of the fittest). The new, stronger pathogenic bacteria are now "resistant" to the established antibiotic, and another antibiotic must be found to fight the new pathogen. We are rapidly approaching that point in history of having super bacteria: disease-causing bacteria that are unaffected by any antibiotic.

The great advantage of Booster® Concentrate is that it does not produce resistant microorganisms during use.

REFERENCES

- Bergsson G, Arnfinnsson J, et al: In vitro inactivation of *Chlamydia trachomatis* by fatty acids and monoglycerides. *Antimicrob Agents Chemother.* 1998 Sep;42(9):2290-4.
- Crouch AA, Seow WK, et al: Effect of human milk and infant milk formulae on adherence of *Giardia intestinalis*. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1991;85:617-619.
- Fletcher RD, Albers AC, et al: Effect of Monoglycerides on *Mycoplasma pneumoniae* Growth. In: *The Pharmacological Effect of Lipids 11*, Kabara JJ (ed) American Oil Chemists' Society: Champaign, Illinois, pp 59-63 (1985).
- Hierholzer JC, Kabara JJ: In Vitro Effects of Monolaurin Compounds on Enveloped RNA and DNA Viruses. *J of Food Safety* 4:1-12 (1982).
- Holland KT, Taylor D, Farrell AM: The effect of glycerol monolaurate on growth of, and production of toxic shock syndrome toxin-1 and lipase by *Staphylococcus aureus*. *J of Antimicrobial Chemo* 1994;33:41-55.
- Homung B, Arntmann E, Sauer G: Lauric acid inhibits the maturation of vesicular stomatitis virus. *J of General Virology* 1994;75:353-361.
- Isaacs CE, Schneidman K: Enveloped Viruses in Human and Bovine Milk are Inactivated by Added Fatty Acids (FAs) and Monoglycerides (MGs). *FASEB Journal* 1991;5: Abstract 5325, p. A1288.
- Isaacs CE, Thormar H: Membrane-disruptive effect of human milk: inactivation of enveloped viruses. *J of Infectious Diseases* 1986; 154:966-971.
- Isaacs CE, Thormar H: The role of milk-derived antimicrobial lipids as antiviral and antibacterial agents. *Adv Exp Med Biol.* 1991;310:159-65.
- Kabara JJ, Ohkawa M, et al: Examinations on Antitumor Immunological, and Plant-Growth Inhibitory Effects of Monoglycerides of Caprylic, Capric, and Lauric Acids and Related Compounds. In: *The Pharmacological Effect of Lipids 11*, Kabara JJ (ed) American Oil Chemists' Society: Champaign, Illinois, pp. 263-272 (1985).
- Kabara JJ (ed): *The Pharmacological Effects of Lipids II*, American Oil Chemists' Society: Champaign, Illinois (1985). Flournoy DJ, Kabara JJ: The Role of Lauricidin[®]E as an Antimicrobial Agent. In: *Drugs of Today* 21(8):373-377 (1985).
- Macallan DC, Noble C, et al: Prospective analysis of patterns of weight change in stage IV human immunodeficiency virus infection. *Am J Clin Nutr.* 1993 Sep;58(3):417-24.
- Oh DH, Marshall DL: Antimicrobial activity of ethanol, glycerol monolaurate or lactic acid against *Listeria monocytogenes*. *Int J Food Microbiol.* 1993 Dec;20(4):239-46.
- Petschow BW, Batema RR, Ford LL: susceptibility of *Helicobacter pylori* to bactericidal properties of medium-chain monoglycerides and free fatty acids. *Antimicrob Agents Chemother.* 1996 Feb;40(2):302-6.
- Projan SJ, Brown-Skrobot S, et al: Glycerol monolaurate inhibits the production of beta-lactamase, toxic shock toxin-1, and other staphylococcal exoproteins by interfering with signal transduction. *J Bacteriol.* 1994 Jul;176(14):4204-9.
- Sands J, Auperin D, Snipes W: Extreme sensitivity of enveloped viruses, including herpes simplex, to long-chain unsaturated monoglycerides and alcohols. *Antimicrob Agents Chemother.* 1979 Jan;15(1):67-73.
- Thormar H, Bergsson G, et al: Hydrogels containing monocaprin have potent microbicidal activities against sexually transmitted viruses and bacteria in vitro. *Sex Transm Infect.* 1999 Jun;75(3):181-5.
- Thormar H, Isaacs CE, et al: Inactivation of enveloped viruses and killing of cells by fatty acids and monoglycerides. *Antimicrob Agents Chemother.* 1987 Jan;31(1):27-31.
- Witcher KJ, Novick RP, Schlievert PM: Modulation of immune cell proliferation by glycerol monolaurate. *Clin and Diagnostic Laboratory Immunology* 1996;3:10-13.